



Math 30 Practice 1st Midterm Solutions

- 1 a) This is exact. It's the same as $(t + \sin(ty) + y^2)' = 0$, so the solution is $t + \sin(ty) + y^2 = C$, which gives y implicitly as a function of t .
- b) This is separable. Either $y=0$, or $\ln(|y|) = t/2 + C$, so $y = D e^{t/2}$.
- c) This is linear, so we find an integrating factor μ . Taking $\mu(t) = e^{t^2}$, we find that $y = \frac{1}{2} + C e^{-t^2}$. Plugging in $y(0) = d$, we get $C = \frac{5}{2}$, so $y = \frac{1}{2} + \frac{5}{2} e^{-t^2}$ is the soln.
- 2) This is linear inhomogeneous. 1st, solve the assoc. homog. problem $y'' + 3y' + dy = 0$. Ans: $y = C_1 e^{-t} + C_2 e^{-dt}$. Next, find a particular soln. Guess: $y = a \cos t + b \sin t$. Solving, we get $a = \frac{1}{10}$, $b = \frac{3}{10}$. So the general soln. is $y = \frac{1}{10} \cos t + \frac{3}{10} \sin t + C_1 e^{-t} + C_2 e^{-dt}$.
- 3) The solns to $y'' - 4y = 0$ are $C_1 e^{2t} + C_2 e^{-2t}$, which tend to 0 iff $C_1 = 0$. Solving $y(0) = y_0$ and $y'(0) = y_0'$ for C_1 , we get $C_1 = \frac{1}{4}(y_0' + dy_0)$, which is 0 if $y_0' = -dy_0$. The solns. for all other initial conditions tend to either $+\infty$ or $-\infty$.
- 4) a) 3rd order, nonautonomous, nonlinear b) 2nd order, autonomous, linear, homogeneous
c) 1st order, autonomous, linear, homogeneous
- 5) It looks like all the solutions fail to exist forever in backward time. If $y_0 = -d$, solns. are not unique.

6) $y' = 0$ when $y = -1$ or 2 :



$y = -1$ is a stable equilibrium, $y = 2$ is unstable. Thus solns. beginning above 2 tend toward $+\infty$, those beginning below -1 tend toward -1 , and those b/w -1 & 2 tend toward -1 .

7) Need to show that t & $t \ln(t)$ are solns (plug in and check), and that the Wronskian is nonzero. $W(t) = \begin{vmatrix} t & t \ln(t) \\ 1 & \ln(t) + 1 \end{vmatrix} = t$, which is $\neq 0$ on $(0, \infty)$, so they do form a fund. set. Set $y = C_1 t + C_2 t \ln(t)$ and plug in $y(1) = 1$, $y'(1) = 2$ to get $y = -t + 3t \ln(t)$.

8) This is separable. Either $y = 0$, or $\frac{1}{y} dy = \frac{1+t}{t} dt \Rightarrow \ln|y| = \ln|t| + t + C$
 $\Rightarrow y = \pm t e^t e^C$, so $y = D t e^t$, where D can be any real number. There are infinitely many solns. with $y(0) = 0$, since $y(0) = 0$ for any choice of D . The theorem does not apply, since in the eqn. $y' = \frac{1+t}{t} y$, $\frac{1+t}{t}$ is not continuous at $t = 0$.

9) Let $A(t) =$ mg in blood at time t . It decays exponentially, so $A(t) = C e^{rt}$ ($A(0) = C$, so C is the initial dose). The half-life is 5 hrs, so $\frac{1}{2} C = C e^{r \cdot 5} \Rightarrow r = -\frac{1}{5} \ln 2$. We want 45.50 mg in the blood at time 1 hr, so $A(1) = C e^{-\frac{1}{5}(\ln 2)} = 45.50 \Rightarrow C = 45.50 \cdot \sqrt[5]{2}$.

10) See section 2.4.